

Aflatoxin B₁ in the Iranian pistachio nut and decontamination methods: A systematic review

Mohsen Mokhtarian^{1*}, Hamid Tavakolipour², Foroud Bagheri³, Carlos Augusto Fernandes Oliveira⁴, Carlos Humberto Corassin⁴, Amin Mousavi Khaneghah^{5,*}

¹Department of Food Science and Technology, Roudehen Branch, Islamic Azad University, Roudehen, Iran;

²Department of Food Science and Technology, Sabzevar Branch, Islamic Azad University, Sabzevar, Iran;

³Department of Food Science and Technology, Yasooj Branch, Islamic Azad University, Yasooj, Iran; ⁴Department of Food Engineering, School of Animal Science and Food Engineering, University of São Paulo, São Paulo, Brazil;

⁵Department of Food Science, Faculty of Food Engineering, University of Campinas (UNICAMP), São Paulo, Brazil

***Corresponding Authors:** Mohsen Mokhtarian, Ph.D., Department of Food Science and Technology, Roudehen Branch, Islamic Azad University, Roudhen, Iran Email: mokhtarian.mo@riau.ac.ir; Amin Mousavi Khaneghah, Ph.D., Department of Food Science, Faculty of Food Engineering, University of Campinas (UNICAMP), Rua Monteiro Lobato, 80. Caixa Postal: 6121.CEP: 13083-862, Campinas, São Paulo, Brazil. Email: mousavi@unicamp.br

Received: 3 July 2020; Accepted: 22 September 2020; Published: 07 October 2020

© 2020 Codon Publications

OPEN ACCESS 

REVIEW ARTICLE

Abstract

This study aimed to summarize the occurrence of aflatoxin B₁ (AFB₁) in the various cultivars of the Persian pistachio nut and the decontamination techniques, with the aid of a systematic review and meta-analysis. In this regard, all published studies up to November 2019 among international and national databases were screened, resulting in 23 articles. According to the findings, the main aflatoxin identified in the pistachio nut was AFB₁, with AFB₁ concentrations in ~28% of the studies being ≤ 5 µg/kg, in ~35% of the studies being ≤ 10 µg/kg, and in ~37% of the studies being ≥ 10 µg/kg. Generally, in most studies (~72%), AFB₁ content in pistachio cultivars was higher than the acceptable level proposed by the Iranian National Standard. Although several physical and chemical techniques for AFB₁ reduction have been introduced, most of them are not safe and/or suitable for human consumption.

Keywords: Pistachio nut; aflatoxin B₁; prevalence; systematic review; decontamination

Introduction

Pistachio (*Pistacia vera* L.), an important plant of the *Anacardiaceae* family, is one of the main Iranian agricultural products (Kola *et al.*, 2018; Mokhtarian *et al.*, 2017; Tavakolipour and Mokhtarian, 2014; Tavakolipour *et al.*, 2017). In 2016, the United States, Iran, and Turkey were the leading pistachio producers globally, reaching 1,335,480, 1,034,999, and 558,303 metric tons, respectively (FAO, 2016). The pistachio nut is consumed in several countries, either as a snack or as an ingredient in food preparation (kernel or powder). A rich composition, including 25% protein (usually essential amino acids),

10% dietary fiber, 55% oil (~80% unsaturated fatty acids), and 16% carbohydrates, provides a high nutritional value for many pistachio-based, traditional Iranian deserts, including Baghlava, Gaz and Ghotab (Mokhtarian *et al.*, 2016; Tavakolipour and Mokhtarian, 2016; Tavakolipour *et al.*, 2017). Although pistachio plays an essential role as a major food item in Iran, more than half of the annual Iranian production is exported, thus requiring appropriate production systems for the exported nuts' quality assurance.

Aspergillus parasiticus, *A. nomius*, and *A. flavus* are important toxigenic fungi that can infect various nuts,

such as pistachio, and contaminate them with aflatoxin G₁ (AFG₁), B₁ (AFB₁), G₂ (AFG₂), and B₂ (AFB₂) (Abdallah *et al.*, 2020; Aydin and Ulvi, 2019; Ehrlich *et al.*, 2007). AFB₁ is a genotoxic carcinogen (Hussein and Brasel, 2001), which leads to toxic effects on the kidney, liver, hematopoietic stem cells, immune system, and the fetal and reproductive systems (Heshmati *et al.*, 2017; Khaneghah *et al.*, 2018, 2019; Nabizadeh *et al.*, 2018). In this regard, the Iranian Scientific Committee for Food (SCF) has reported that AFB₁ is a major factor in cancer, especially for the liver, even at very low amounts (Bensassi *et al.*, 2010). According to Iran's national standard, the maximum tolerated levels for AFB₁ and the total aflatoxin (TAFT) in pistachio are 5 and 15 µg/kg, respectively (ISIRI, 2002). However, the European Commission standardized the aflatoxin (AFT) content of pistachio as <10 µg/kg (European Commission, 2010). The prevalence of AFT in the pistachio nuts has been reported previously. In 2015, the Rapid Alert System for Food and Feed (RASFF) reported a total of 475 alerts for mycotoxins, of which 89% were associated with AFT, mostly from nuts (81%), mainly pistachio nuts, pumpkin seeds, and dried mix nuts (Pigłowski, 2019; RASFF, 2016). Georgiadou *et al.* (2012) investigated the AFT contamination in the pistachio nuts. In order to determine the rate of AFT contamination in the pistachio nuts, four orchards were chosen, and 20 samples (total of 80 samples) were selected from each orchard. It should be noted that AFT control of food commodities requires well-developed analytical methods (Mousavi Khaneghah *et al.*, 2018). The combination and structure of food products, identification, and mycotoxins detection is a challenge (Campagnollo *et al.*, 2016; Heshmati *et al.*, 2017). Therefore, analytical methods for precise quantification of mycotoxins in the pistachio nuts have been developed, aiming to provide high sensitivity, easy-to-operate steps, rapidness, and low-cost sample preparation procedures (Rahmani *et al.*, 2018).

In recent years, several technological approaches have been introduced to reduce the growth of fungi or for the removal of AFT from pistachios, including the application of extracts and essential oils from medicinal plants (Ijabadeniyi *et al.*, 2020; Khorasani *et al.*, 2017); ozone treatment (O₃) (Chen *et al.*, 2014); the use of chemical compounds, such as methyl bromide (CH₃Br), hydrogen peroxide (H₂O₂), sodium bisulfite (NaHSO₃), and ammonia; the use of adsorbent compounds, such as active charcoal, zeolite, sodium bentonite, hydrated sodium calcium aluminosilicate (HSCAS); gamma radiation (Serra *et al.*, 2018); and biological methods, such as mannan compounds from the cell wall of *Saccharomyces cerevisiae* and lactic acid bacteria (Abdallah *et al.*, 2018; Abdolshahi *et al.*, 2018). Examples of the application of biological methods include the report by Abdolshahi *et al.* (2018),

who studied the interacting capacity of *S. cerevisiae* mannanoprotein (mannan) with AFT in pistachio by soaking the kernels in mannan solutions at 25 and 50 mg/mL. Siahmoshteh *et al.* (2017) investigated the efficacy of *B. amyloliquefaciens* and *B. subtilis* in controlling AFT production and *Aspergillus parasiticus* growth. The authors considered the *Bacillus* species as good biocontrol agents to reduce toxigenic fungal material growth and the subsequent AFT decontamination of pistachio. Rastegar *et al.* (2017) investigated the removal of AFB₁ by roasting contaminated pistachio nuts mixed with lemon juice and/or citric acid.

It can be stated that, due to the economic, nutritional, and health aspects of the Iranian pistachio nut (Mokhtarian, 2015; Ros, 2010), along with the high susceptibility of this product to fungal growth (especially to *Aspergillus* species), there is a need for strategies to reduce AFT contamination at different stages of harvest, postharvest, and during the processing of this product. Therefore, this study aimed to review the data from all available studies on the prevalence of AFB₁ in different Iranian pistachio cultivars and recommend effective ways to combat this severe challenge.

Search Strategy

Keywords and screened databases

In order to study the prevalence of AFB₁ in the different Iranian pistachio cultivars, all published studies up to December of 2018 were screened using "Scopus," "PubMed," and "SID" databases using the keywords, "Pistachio nut," "Pistachio kernel," "Aflatoxin contamination," "*Aspergillus flavus*," "Anti-aflatoxigenic," "Anti-mycotoxigenic," "Iranian," and "Persian." Also, all references in the retrieved studies were reviewed to prevent any study from being missed out. It should be noted that all studies that examined the analytical methods of AFT detection in pistachios were excluded from the research.

Data extraction

Considering that the pistachio is one of Iran's nonoil and strategic products, in this study, only the research carried out by Iranian researchers was addressed. From each study, the required information, including author name (researcher), publication year, analytical technique, pistachio cultivar, an average of the reported AFB₁, number of samples, the ratio of the measured AFT to the maximum tolerated level of AFB₁ following the Iranian national standard AFB_{1R}/AFB_{1S}, and the method applied to reduce AFTs in pistachio were obtained.

Main Findings

Study characteristics

In the initial evaluation of the 1582 articles collected, 674 were excluded because of duplication (repeated articles) and irrelevant data descriptions. A total of 908 articles extracted as the primary articles based on abstracts and

titles were excluded because 835 articles reported on other toxins and 73 articles did not describe the analytical method used. Only those researches reporting the AFT contamination with the particular analytical method used and studies on decontamination techniques in pistachio nuts were evaluated. At the end of the data extraction procedure, 23 articles published before 30 December 2018 were selected and evaluated in our study (Tables 1, 2, and Figure 1).

Table 1. Published researches related to the prevalence of AFB₁ contamination in the pistachio nuts produced in Iran.

Pistachio cultivar	Average of AFB ₁ reported in research (µg/kg)	Number of studied samples (N)	Ratio $\frac{AFB_{1R}^{**}}{AFB_{1S}^{***}}$	Identification method	Ref.
O'hadi	285.5	10	~57	HPLC	Mehrnezhad and Panahi, 2005
	1485	14	~297		
O'hadi	6.25	–	1.25	TLC	Pak-kish and Rahemi, 2005
Akbari	10	–	2	HPLC	Rahaie <i>et al.</i> , 2010b
Akbari	5.63	–	1.126	HPLC	Tavakolipour <i>et al.</i> , 2012
Fandoghi (Early split)	–	7.93	~1.59	ELISA	Dargahi <i>et al.</i> , 2014
Fandoghi (Whole)	1.59	–	~0.32	ELISA	Dargahi <i>et al.</i> , 2014
Unknown	~0.681	72	~0.136	HPLC	Shakeri and Fazaeli, 2016
Kermani	5.9	3356	1.18	HPLC	Cheraghali <i>et al.</i> , 2007
Unknown	8.12	–	~1.63	HPLC	Aghamohammadi <i>et al.</i> , 2007
Kermani	~311.6	3181	~62.32	HPLC	Dini <i>et al.</i> , 2013
Ahmad Aghaei (Damghan)	2.077	600	~0.42	HPLC	Taghizadeh <i>et al.</i> , 2018
Akbari (Damghan)	1.948	600	0.3896		
Kalle-Ghuchi (Damghan)	1.627	600	0.3254		
O'hadi (Damghan)	<0.066	600	0.0132		
Badami (Damghan)	<0.066	600	0.0132		
Ahmad Aghaei (Feyz Abad)	3.863	600	0.7726		
Akbari (Feyz Abad)	3.081	600	0.6162		
Kalle-Ghuchi (Feyz Abad)	2.143	600	0.4286		
O'hadi (Feyz Abad)	0.2	600	0.04		
Badami (Feyz Abad)	0.2	600	0.04		
Ahmad Aghaei (Rafsanjan)	4.33	600	0.866		
Akbari (Rafsanjan)	4.08	600	0.816		
Kalle-Ghuchi (Rafsanjan)	3.453	600	0.6906		
O'hadi (Rafsanjan)	0.2	600	0.04		
Badami (Rafsanjan)	0.2	600	0.04		
Ahmad Aghaei (Sarakhs)	0.961	600	0.1922		
Akbari (Sarakhs)	0.487	600	0.0974		
Kalle-Ghuchi (Sarakhs)	0.233	600	0.0466		
O'hadi (Sarakhs)	<0.066	600	0.0132		
Badami (Sarakhs)	<0.066	600	0.0132		
Unknown	22.02****	43	~4.4	ELISA	Ostadrahimi <i>et al.</i> , 2014

*Maximum tolerated level of AFB₁ in accordance with the Iranian National Standard (No. 5925, 1380), 5 µg/kg (ISIRI, 2002).

**AFB_{1R}: Obtained AFB₁ content in accordance with the research.

***AFB_{1S}: The AFB₁ content in accordance with the Iranian National Standard (1380).

****In this research, the noted value is for TAFT.

AFBB₁, aflatoxin B₁; HPLC, high-performance liquid chromatography; TLC, thin-layer chromatography; ELISA, enzyme-linked immunosorbent assay.

Table 2. The published applied researches related to the decontamination methods of AFB₁ in the pistachio nuts produced in Iran.

Pistachio cultivar	Average of AFB ₁ (µg/kg)		Aflatoxin decontamination method	Amount of toxin removal (%)	Ref.
	Before treatment	After treatment			
O'hadi	6.25	1.2	25 mg/l Gibberellic acid	~81	Pak-kish and Rahemi, 2005
Ahmad Aghaei	~2.75	~0.5	Phenolics of pistachio green hull	~81	Afshari et al., 2009
Akbari	10	6	Inoculation of pistachio by <i>Saccharomyces cerevisiae</i> yeast	40	Rahaie et al., 2010a
Akbari	16.467	5.644	Extract of Shirazi thyme	~66	Tavakolipour et al., 2010
Shahpasand	32,300	31,171	Testa (Pistachio kernel coat)	~3.49	Mohammadi Moghadam et al., 2011
Abasali	32,307	30,754		~4.81	
O'hadi	32,810	29,823		~9.11	
Ahmad Aghaei	30,171	29,429		~2.46	
Akbari	29,940	26,880		~10.22	
Kale-Bozi	27,439	26,410		~3.75	
Kalle-Ghuchi	30,433	26,218		~13.85	
FAS-13-73	29,012	25,393		~12.47	
Fakhri	30,762	24,811		~19.35	
Kal-e-Khandan	27,912	21,728		~22.15	
Akbari	16.467	5.644	Extract of the Shirazi thyme	~66	Tavakolipour et al., 2012
Unknown	1700.63	610.32	<i>Trichoderma</i> spp. T ₁ extracellular extract	~64.11	Chegini, Behbodi et al., 2013
		265.29	<i>Trichoderma</i> spp. T ₃ extracellular extract	~84.4	
		376.03	<i>Trichoderma</i> spp. T ₄ extracellular extract	~77.88	
		825.75	<i>Trichoderma</i> spp. T ₁₇ extracellular extract	~51.44	
O'hadi	–	–	4 ppm soluble ozone in water	32.7	Bashiri et al., 2013
	–	–	8 ppm soluble ozone in water	47.9	
Unknown	–	–	Radiation at 1 kGy	38.84	Haji Mohammadi et al., 2016
	–	–	Radiation at 3 kGy	48.79	
	–	–	Radiation at 5 kGy	53.50	
	–	–	Radiation at 7 kGy	77.17	
Unknown (Pistachio kernel)	44		Roasted at 150°C (30 min)	66	Yazdanpanah et al., 2005
	91			64	
	213	–		24	
Unknown (Whole pistachio)	144	–	Roasted at 150°C (30 min)	63	
	253	–		19	
Unknown	~2500	~450	Inoculation of pistachio by <i>Bacillus subtilis</i> (UTBSP1) for 96 h at 30°C	82	Farzaneh et al., 2012
Ahmad Aghaei	–	–	Inoculation of pistachio by <i>Bacillus amyloliquefaciens</i> (UTBSP1) for 5 days at 30°C	23.9	Siahmoshteh et al., 2017
	–	–	Inoculation of pistachio by <i>Bacillus amyloliquefaciens</i> (UTBSP1) for 8 days at 30°C	54.9	
	–	–	Inoculation of pistachio by <i>Bacillus subtilis</i> for 5 days at 30°C	41.1	
	–	–	Inoculation of pistachio by <i>Bacillus subtilis</i> for 8 days at 30°C	52.5	

(continues)

Table 2. Continued

Pistachio cultivar	Average of AFB ₁ (µg/kg)		Aflatoxin decontamination method	Amount of toxin removal (%)	Ref.
	Before treatment	After treatment			
Unknown	268	–	Roasting of pistachio at 120°C for 60 min by incorporation of 30 mL distilled water, 15 mL lemon juice, 2.25 g citric acid, and 5 g NaCl	58.6	Rastegar <i>et al.</i> , 2017
	268	–	Roasting of pistachio at 150°C for 60 min by incorporation of 30 mL distilled water, 15 mL lemon juice, 2.25 g citric acid, and 5 g NaCl	47	
Unknown	–	–	Soaking of pistachio in mannan solution (25 mg/mL)	63.4	Abdolshahi <i>et al.</i> , 2018
	–	–	Soaking of pistachio in mannan solution (50 mg/mL)	84.4	
Unknown	~2500	~1750	Inoculation of pistachio by <i>Bacillus subtilis</i> (UTB1) for 72 h at 37°C	30	Afsharmanesh <i>et al.</i> , 2018
	~2500	~1750	Inoculation of pistachio by <i>Bacillus subtilis</i> (UTB1) and mutant (B1Δ <i>bacC5</i>) for 72 h at 37°C	30	
	~2500	~1580	Inoculation of pistachio by <i>Bacillus subtilis</i> (UTB1) and mutant (B1Δ <i>bacC12</i>) for 72 h at 37°C	36.8	
	~2500	~1000	Inoculation of pistachio by <i>Bacillus subtilis</i> (UTB1) and mutant mutant (B1Δ <i>bacC18</i>) for 72 h at 37°C	60	
Akbari	10	–	Inoculation of pistachio by <i>Saccharomyces cerevisiae</i> yeast for 12 h at 25°C	40	Rahaie <i>et al.</i> , 2010a
	20	–		70	
Akbari	10	–	Inoculation of pistachio by <i>Saccharomyces cerevisiae</i> yeast for 12 h at 25°C along with acidic treatment (2M HCL)	60	
	20	–		73	
Akbari	10	–	Inoculation of pistachio by <i>Saccharomyces cerevisiae</i> yeast for 12 h at 25°C along with heat treatment (autoclave at 121°C for 20 min)	55	
	20	–		75	
Akbari	–	–	Addition of Salicylic acid solution (1 mmol/l) to medium culture	~30	Panahirad <i>et al.</i> , 2014
	–	–	Addition of Salicylic acid solution (3 mmol/l) to medium culture	~42	
	–	–	Addition of Salicylic acid solution (5 mmol/l) to medium culture	~50	
	–	–	Addition of Salicylic acid solution (7 mmol/l) to medium culture	~87	
	–	–	Addition of Salicylic acid solution (9 mmol/l) to medium culture	~100	
Akbari	–	–	Addition of Salicylic acid solution (11 mmol/l) to medium culture	~100	Panahirad <i>et al.</i> , 2014
Unknown	22.02**	0.48	Roasting of pistachio with salt	~98	Ostadrhimi <i>et al.</i> , 2014

*Maximum tolerated level of AFB, following the Iranian National Standard (No. 5925, 1380), 5 µg/kg (ISIRI, 2002).

**In this research, the noted value is for TAFT AFB₁, aflatoxin B₁.




Figure 1. Flow chart of exclusion and inclusion studies based on the PRISMA guidelines.

Prevalence of AFB₁ in the Iranian pistachio

According to Table 1, the evaluation and measurement of AFB₁ content in the Iranian pistachio has been described in only 11 studies. Our work results indicate that the Iranian pistachio cultivars evaluated for AFT content include O'hadi, Akbari, Kermani, Ahmad Aghaei, Kalle-Ghuchi, and Badami. Generally, the AFB₁ concentrations of various Persian pistachio cultivars varied from <0.066 to 1,485 µg/kg (Table 1). The results demonstrated that, with time, the AFT content has decreased, probably because of increasing awareness among farmers concerning improving the sanitary conditions in the orchards, vehicles, and pistachio terminals. As per the results, among the published articles on the incidence of AFT in the Persian pistachio, only two studies reported AFB₁ content higher than 285.5 µg/kg, while in the other articles, the AFB₁ levels varied markedly from 0.066 to 10 µg/kg ($0.066 \leq \text{AFB}_1 (\mu\text{g}/\text{kg}) \leq 10$). The AFT levels in ~28% of the studies were lower than 5 µg/kg ($\text{AFB}_1 \leq 5 \mu\text{g}/\text{kg}$), while ~35% of the studies reported concentrations in the range of $5 \leq \text{AFB}_1 (\mu\text{g}/\text{kg}) \leq 10$ and ~37% of the studies mentioned levels $> 10 \mu\text{g}/\text{kg}$ ($\text{AFB}_1 \geq 10 \mu\text{g}/\text{kg}$). As can be seen, the AFB₁ content in a significant number of studies (~72%) was higher than the acceptable limit range set by the national standard of Iran (i.e., < 5 µg/kg).

Georgiadou *et al.* (2012) reported that a critical point for AFT contamination of the pistachio nuts is the

maturity step, ranging from 11 to 1,361 µg/kg among orchards. In the harvesting step, the AFT content was >1,420 µg/kg, and the authors concluded that the higher AFT levels (>1000 µg/kg) in orchards correlated with the highest insect infestations during this production step. At the postharvest level, AFT contamination in three out of four orchards varied from 650 to 1,100 µg/kg. During the storage period, it ranged between 40 and 1,200 µg/kg.

Considering that some environmental factors are needed to support fungal growth, such as temperature and the relative humidity of the environment and storage, proper drying and suitable storage of grains are the most important preventive actions to prevent fungal growth. In this context, fungal contamination and the subsequent AFT production in pistachio nuts may occur in orchards and farms, at harvesting time, during transportation, during the process stage, and storage, and as well as when distributing the product (Cheraghali *et al.*, 2007).

In this study, 23 articles were found related to AFT's prevalence in the Iranian pistachio, which indicated that the most frequent AFT found in pistachio is AFB₁. Several studies have confirmed the presence of AFB₁ in the Iranian pistachio (Afsharmanesh *et al.*, 2018; Panahirad *et al.*, 2014; Rahaie *et al.*, 2010a; Yazdanpanah *et al.*, 2005). Dini *et al.* (2013) studied the AFT contamination levels in the Iranian pistachio and concluded that

the mean AFB₁ levels in this product in 2009, 2010, and 2011 were ~320 246, and ~370 µg/kg, respectively. They also observed that the measured AFT was higher than those recommended by the Iranian National Standard in all the years mentioned above. In another study, Cheraghali et al. (2007) claimed that AFB₁ content in ~63.3% of the pistachio samples was lower than the limit of detection (LOD) of the analytical method used in the study (5 µg/kg).

Analytical methods for the determination of AFT

Today, the successful development of fast, robust, cheap, and simple analytical methods with high selectivity and sensitivity for measuring and identifying different mycotoxins in different foods products is a reality (Alshannaq and Yu, 2017; Zhang et al., 2018). New methods rely on gas and high-performance liquid chromatography (HPLC) systems combined with tandem mass spectrometry (MS/MS), thereby providing an unequivocal identification and precise quantification of multiple mycotoxins in complex food matrices (Turner et al., 2015). Also, HPLC coupled

with fluorescence detection (FLD) was developed as one of the nonMS methods earlier by the European Standardization Committee (CEN) and the Association of Official Analytical Chemists (AOAC) International for the determination and identification of different mycotoxins (Kwaśniewska et al., 2015; Pascale, 2009; Pereira et al., 2014). The HPLC-FLD sensitivity is comparable to the results determined by the HPLC-MS/MS technique, but the HPLC-FLD is usually more suitable for single mycotoxin detection or the determination of mycotoxins group with similar chemical structures (Khaneghah et al., 2019). In all the literature studied (Figure 2a), the analytical methods based on HPLC were the most used for the determination of AFTs in the pistachio nuts, followed by thin-layer chromatography (TLC) and enzyme-linked immunosorbent assay (ELISA).

The ELISA method is based on the chemical interaction between a given antigen (in this case, a mycotoxin conjugated to the larger, immunogenic molecule) and its specific antibody. According to this interaction arrangement, ELISA protocols can be designed as direct, indirect, sandwich ELISA, competitive, or indirect competitive. In the




Figure 2. The number of articles and studies on identification methods (A), year of publishing (D), the decontamination method (C), and the pistachio type (D).

direct ELISA, the extract of samples reacts directly with the enzyme, thus providing rapid and beneficial results for mycotoxins' qualitative evaluation. Competitive ELISA shows the competitive reaction between sample (antibody or antigen) and target mycotoxin in the enzyme-labeled targets versus immobilized antibody or antigen. The indirect technique is based on indirectly determined target antigen using the secondary antibody, which is usually labeled as an enzyme. The indirect competitive technique's mechanism is the interaction of the primary antibody with the immobilized antigen, and in this method, a secondary antibody conjugated with an enzyme is required. In the sandwich technique, the antigen interacts with two different antibodies (Suzuki *et al.*, 2007), which allows high specificity, rapid speed screening, simple operation without using any readers or instrument, and eco-friendly analysis due to radioactive labels and low volumes of solvents (Batrinou *et al.*, 2020; Krska and Molinelli, 2009; Sakamoto *et al.*, 2018).

One of the most critical limitations of the regular monitoring of mycotoxins in foods using ELISA is the high price of commercially available kits, especially in developing countries (Rossi *et al.*, 2012). The further investments in immunoreagent manufacturing could increase the access to the kits as ELISA is the most common immunoassay applied for fast mycotoxin determination. Other fast immunoassay-based methods, including radioimmunoassay, immunoaffinity columns, and sequential injection immunoassay, have been applied to identify and determine mycotoxins in food products. Moreover, rapid, on-site test bands have been developed to measure toxic contaminants in foodstuffs, including a flow-through, lateral-flow immunoassay (LFIA), and dipstick. The lateral flow-based assay is a big, sensitive, selective, and single-step test for determining different mycotoxins in about 10 min. This technique is based on three factors: a porous membrane, an absorbent pad, and a conjugate pad. These three factors ensure a lateral flow, increase the flowing liquid volume, and guarantee contact between the liquid sample and the membrane (Maragos and Busman, 2010).

Methods for the decontamination of AFTs in the Iranian pistachio

The experimental techniques applied for AFT decontamination of the pistachio are presented in Table 2. It is noteworthy that the majority of the studies were conducted using Iranian pistachio cultivars such as O'hadi, Ahmad Aghaei, Akbari, Shahpasand, Abasali, Kale-Bozi, Kalle-Ghuchi, FAS-13-73, Fakhri, and Kal-e-Khandan (Table 2, Figure 2b and d). In accordance to the results reported in the evaluated studies, the lowest (2.46%) and the highest (100%) percentages of AFB₁ removal were related to

treatments using "pistachio Testa (pistachio kernel coat)" and "addition of 9 mmol/l Salicylic acid solution to the medium culture," respectively. Overall, the applied methods for the decontamination of AFTs in the Iranian pistachio included physical (radiation and roasting) and chemical methods (gibberellic acid, treatment with ozone, lemon extract, citric acid, salt solution, and salicylic acid solution). While biological treatments, such as *Saccharomyces cerevisiae*, *Trichoderma* extracellular extract, inoculation by *Bacillus* species, and recombinant mutants of *Bacillus subtilis*, attracted notable attention, combined methods and the use of extract and essential oils from medical plants, such as phenolic compounds from pistachio green hull, the extract of the Shirazi thyme, and Testa or the pistachio kernel coat, were also useful (Figure 2c).

Abdolshahi *et al.* (2018) studied the AFT-binding efficiency of *S. cerevisiae* mannoprotein (mannan) in contaminated pistachio nuts, observing the highest binding percentage (84.4%) for AFB₁ at a higher concentration of mannan (50 mg/mL). In another research, the simultaneous application of roasting-chemical treatment (lemon juice and citric acid) for the reduction of AFB₁ in pistachio was evaluated by Rastegar *et al.* (2017). The authors found that roasting of pistachio nuts (50 g) in water (30 mL), lemon juice (30 mL), and citric acid (6 g) at 120°C for 60 min resulted in a significant reduction (93.1%) of AFB₁. Also, the roasting of pistachio nuts (50 g) with water (30 mL), lemon juice (15 mL), and citric acid (2.25 g) at 120 C for 60 min decreased the level of AFB₁ to 49.2% of the original content. Generally, it was reported that the simultaneous use of heat and citric acid/lemon juice had a synergistic effect on AFB₁ reduction. In summary, physical, chemical, and biological methods, as proposed in several studies, could significantly reduce the pistachio nuts' AFT contents. However, methods based on the inoculation of different microorganisms that are generally recognized as safe (GRAS), such as *B. subtilis* and *S. cerevisiae*, have a higher potential for practical application pistachio industry. Siahmoshteh *et al.* (2017) demonstrated that both strains could decrease the mycelial growth *in vitro*, reducing the production of AFB₁, AFB₂, AFG₁, and AFG₂ during the first 3 days after inoculation.

Conclusions

The results of this study indicate that the occurrence levels of AFB₁ in the Persian pistachio cultivars are a public health concern as 72% of the studies reported concentrations higher than the acceptable level per the Iranian National Standard for this product. The high incidence of AFTs in the Iranian pistachio is probably due to a lack of knowledge among the farmers, pistachio terminals, and the pistachio processing stages on the preventive

measures with regard to toxigenic fungi contamination of foodstuffs. Therefore, further preventive actions and also ongoing monitoring can counter the potential risk of AFT in pistachios. Although several experimental techniques have been proposed for the reduction of the AFT content in pistachio nuts, the inoculation of GRAS microorganisms shows the most significant potential for future applications in the pistachio industry.

References

- Abdallah, M.F., Ameye, M., De Saeger, S., Audenaert, K. and Haesaert, G., 2018. Biological control of mycotoxigenic fungi and their toxins: An update for the pre-harvest approach. In: Mycotoxins-impact and management strategies. IntechOpen. Patrick Berka Njobeh and Francois Stepman, London, UK, <https://doi.org/10.5772/intechopen.76342>
- Abdallah, M.F., Audenaert, K., Lust, L., Landschoot, S., Bekaert, B., Haesaert, G., De Boevre, M. and De Saeger, S., 2020. Risk characterization and quantification of mycotoxins and their producing fungi in sugarcane juice: A neglected problem in a widely-consumed traditional beverage. *Food Control* 108: 106811. <https://doi.org/10.1016/j.foodcont.2019.106811>
- Abdolshahi, A., Shabani, A.A., Mortazavi, S.A. and Marvdashti, L.M., 2018. Aflatoxin binding efficiency of *Saccharomyces cerevisiae* mannoprotein in contaminated pistachio nuts. *Food Control* 87: 17–21. <https://doi.org/10.1016/j.foodcont.2017.12.008>
- Afshari, H., Talaie, A., Mohammadi Moghadam, M. and Panahi, B., 2009. Differences of elements in early splitting of pistachio nuts and the effect of phenolic compounds and gallic acid on rate. *Journal of Horticulture Science* 23: 10–17. [In Persian]
- Afsharmanesh, H., Perez-Garcia, A., Zerriouh, H., Ahmadzadeh, M. and Romero, D., 2018. Aflatoxin degradation by *Bacillus subtilis* UTB1 is based on production of an oxidoreductase involved in bacilysin biosynthesis. *Food Control* 94: 48–55. <https://doi.org/10.1016/j.foodcont.2018.03.002>
- Aghamohammadi, M., Hashemi, J., Kram, G.A. and Alizadeh, N., 2007. Enhanced synchronous spectrofluorimetric determination of aflatoxin B₁ in pistachio samples using multivariate analysis. *Analytica Chimica Acta* 582: 288–294. <https://doi.org/10.1016/j.aca.2006.09.014>
- Alshannaq, A. and Yu, J.-H., 2017. Occurrence, toxicity, and analysis of major mycotoxins in food. *International Journal of Environmental Research and Public Health* 14: 632. <https://doi.org/10.3390/ijerph14060632>
- Aydin, S. and Ulvi, M., 2019. Residue levels of pesticides in nuts and risk assessment for consumers. *Quality Assurance and Safety of Crops & Foods* 11: 539–548. <https://doi.org/10.3920/QAS2018.1405>
- Bashiri, P., Hadad Khodaparast, M.H., Sedaghat, N., Tabatabaei Yazdi, F. and Nasiri Mohalati, M., 2013. The effects of solution ozone in water on aflatoxin reduction of O'hadi pistachio cultivar. *Iranian Food Science and Technology Research Journal* 9: 215–221. [In Persian]
- Batrinou, A., Houhoula, D. and Papageorgiou, E., 2020. Rapid detection of mycotoxins on foods and beverages with enzyme linked immunosorbent assay. *Quality Assurance and Safety of Crops & Foods* 12: 40–49. <https://doi.org/10.15586/QAS2019.654>
- Bensassi, F., Rhouma, A., Ghrab, M., Bacha, H. and Hajlaoui, M.R., 2010. Evaluation of cultivar susceptibility and storage periods towards aflatoxin B₁ contamination on pistachio nuts. *Mycotoxin Research* 26: 199–203. <https://doi.org/10.1007/s12550-010-0054-7>
- Campagnollo, F.B., Ganey, K.C., Khaneghah, A.M., Portela, J.B., Cruz, A.G., Granato, D., Corassin, C.H., Oliveira, C.A.F. and Sant'Ana, A.S., 2016. The occurrence and effect of unit operations for dairy products processing on the fate of aflatoxin M₁: A review. *Food Control* 68: 310–329. <https://doi.org/10.1016/j.foodcont.2016.04.007>
- Chegini, S., Behbodi, K., Javan Nikkhah, M. and Farzaneh, M., 2013. Study of the effects of *Trichoderma* isolates on prevent of fungal growth of *Aspergillus flavus* and aflatoxin reduction on pistachio. *Journal of Biological Control Pests Plant Diseases* 2: 71–79. [In Persian]
- Chen, R., Ma, F., Li, P.-W., Zhang, W., Ding, X.-X., Zhang, Q., Li, M., Wang, Y.-R. and Xu, B.-C., 2014. Effect of ozone on aflatoxins detoxification and nutritional quality of peanuts. *Food Chemistry* 146: 284–288. <https://doi.org/10.1016/j.foodchem.2013.09.059>
- Cheraghali, A., Yazdanpanah, H., Doraki, N., Abouhossain, G., Hassibi, M., Ali-Abadi, S., Aliakbarpoor, M., Amirahmadi, M., Askarian, A. and Fallah, N., 2007. Incidence of aflatoxins in Iran pistachio nuts. *Food and Chemical Toxicology* 45: 812–816. <https://doi.org/10.1016/j.fct.2006.10.026>
- Dargahi, R., Moradi, M., Fani, S.R. and Masoumi, H., 2014. Evaluation of aflatoxin B₁ in different parts of pistachio fruit and effects of processing stages. *Journal of Food Hygiene* 4: 21–31. [In Persian]
- Dini, A., Khazaeli, P., Roohbakhsh, A., Madadlou, A., Pourenamdari, M., Setoodeh, L., Askarian, A., Doraki, N., Farrokhi, H. and Moradi, H., 2013. Aflatoxin contamination level in Iran's pistachio nut during the years 2009–2011. *Food Control* 30: 540–544. <https://doi.org/10.1016/j.foodcont.2012.08.012>
- Ehrlich, K.C., Kobbeman, K., Montalbano, B.G. and Cotty, P.J., 2007. Aflatoxin-producing *Aspergillus* species from Thailand. *International Journal of Food Microbiology* 114: 153–159. <https://doi.org/10.1016/j.ijfoodmicro.2006.08.007>
- European Commission, 2010. Amending regulation setting maximum levels, for certain contaminants in foodstuffs as regards aflatoxins. *Official Journal of the European Union* 1881- L50: 8–12.
- FAO, 2016. FAOSTAT-Statistical Service, FAO, Rome.
- Farzaneh, M., Shi, Z.-Q., Ghassempour, A., Sedaghat, N., Ahmadzadeh, M., Mirabolfathy, M. and Javan-Nikkhah, M., 2012. Aflatoxin B₁ degradation by *Bacillus subtilis* UTBSP1 isolated from pistachio nuts of Iran. *Food Control* 23: 100–106. <https://doi.org/10.1016/j.foodcont.2011.06.018>
- Georgiadou, M., Dimou, A. and Yanniotis, S., 2012. Aflatoxin contamination in pistachio nuts: A farm to storage study. *Food Control* 26: 580–586. <https://doi.org/10.1016/j.foodcont.2012.02.014>

- Haji Mohammadi, B., Ehrampoush, M.H., Hashemi, S., Khalatbari Limaki, S., Zare, F. and Taheri Soudejani, M., 2016. The effect of electron irradiation on aflatoxin B₁ in pistachio production process inoculated with *Aspergillus flavus*. *Journal of Toloo-e-Behdasht* 16: 1–8. [In Persian]
- Heshmati, A., Zohrevand, T., Khaneghah, A.M., Mozaffari Nejad, A.S. and Sant'Ana, A.S., 2017. Co-occurrence of aflatoxins and ochratoxin A in dried fruits in Iran: Dietary exposure risk assessment. *Food and Chemical Toxicology* 106: 202–208. <https://doi.org/10.1016/j.fct.2017.05.046>
- Hussein, H.S. and Brasel, J.M., 2001. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology* 167: 101–134. [https://doi.org/10.1016/S0300-483X\(01\)00471-1](https://doi.org/10.1016/S0300-483X(01)00471-1)
- Ijabadeniyi, O., Mbedla, A. and Ajayeoba, T., 2020. Microbiological quality and antimicrobial efficacy of combined oregano essential oil and acetic acid on fresh lettuce. *Italian Journal of Food Science* 32(2): 399–409. <https://doi.org/10.14674/IJFS-1566>
- ISIRI (Institute of Standard and Industrial Research of Iran), 2002. Maximum tolerated limits of mycotoxins in foods and feeds. National Standard No.5925. ISIRI, Karaj, Iran.
- Khaneghah, A.M., Fakhri, Y., Gahrue, H.H., Niakousari, M. and Sant'Ana, A.S., 2019. Mycotoxins in cereal-based products during 24 years (1983–2017): A global systematic review. *Trends in Food Science & Technology* 91: 95–105. <https://doi.org/10.1016/j.tifs.2019.06.007>
- Khaneghah, A.M., Fakhri, Y., Raeisi, S., Armoon, B. and Sant'Ana, A.S., 2018. Prevalence and concentration of ochratoxin A, zearalenone, deoxynivalenol and total aflatoxin in cereal-based products: A systematic review and meta-analysis. *Food and Chemical Toxicology* 118: 830–848. <https://doi.org/10.1016/j.fct.2018.06.037>
- Khorasani, S., Azizi, M.H., Barzegar, M., Hamidi-Esfahani, Z. and Kalbasi-Ashtari, A., 2017. Inhibitory effects of cinnamon, clove and celak extracts on growth of *Aspergillus flavus* and its aflatoxins after spraying on pistachio nuts before cold storage. *Journal of Food Safety* 37: e12383. <https://doi.org/10.1111/jfs.12383>
- Kola, O., Hayoğlu, İ., Türkoğlu, H., Parıldı, E., Ak, B.E. and Akkaya, M.R., 2018. Physical and chemical properties of some pistachio varieties (*Pistacia vera* L.) and oils grown under irrigated and non-irrigated conditions in Turkey. *Quality Assurance and Safety of Crops & Foods* 10: 383–388. <https://doi.org/10.3920/QAS2017.1152>
- Krska, R. and Molinelli, A., 2009. Rapid test strips for analysis of mycotoxins in food and feed. *Analytical and Bioanalytical Chemistry* 393: 67–71. <https://doi.org/10.1007/s00216-008-2424-y>
- Kwaśniewska, K., Gadzała-Kopciuch, R. and Cendrowski, K., 2015. Analytical procedure for the determination of zearalenone in environmental and biological samples. *Critical Reviews in Analytical Chemistry* 45: 119–130. <https://doi.org/10.1080/10408347.2014.896731>
- Maragos, C. and Busman, M., 2010. Rapid and advanced tools for mycotoxin analysis: A review. *Food Additives and Contaminants* 27: 688–700. <https://doi.org/10.1080/19440040903515934>
- Mehrnezhad, M.R. and Panahi, B., 2005. The influence of hull cracking on aflatoxin contamination and insect infestation in pistachio nuts. *Entomology and Phytopathology* 73: 105–123. [In Persian]
- Mohammadi Moghadam, M., Afshari, H., Mohammadi Goltape, E., Hokmabadi, H. and Rad, S., 2011. Investigation of the effects of testa in reducing growth of *Aspergillus flavus* fungi aflatoxin and production of aflatoxin B₁ in different cultivars of pistachio kernel. *Journal of Horticulture Science* 25: 73–81. [In Persian]
- Mokhtarian, M., 2015. Thermodynamic study of a solar dryer equipped with air recycle and uses it to modeling of pistachio drying process by using genetic algorithm and artificial neural network approaches. PhD dissertation. Islamic Azad University, pp. 1–205. [In Persian]
- Mokhtarian, M., Tavakolipour, H. and Ashtari, A.K., 2017. Effects of solar drying along with air recycling system on physicochemical and sensory properties of dehydrated pistachio nuts. *LWT* 75: 202–209. <https://doi.org/10.1016/j.lwt.2016.08.056>
- Mokhtarian, M., Tavakolipour, H. and Kalbasi-Ashtari, A., 2016. Energy and exergy analysis in solar drying of pistachio with air recycling system. *Drying Technology* 34: 1484–1500. <https://doi.org/10.1080/07373937.2015.1129499>
- Mortazavi, A. and Tabatabai, F., 1998. Mycotoxins, pp. 1–206. [In Persian]
- Mousavi Khaneghah, A., Fakhri, Y. and Sant'Ana, A.S., 2018. Impact of unit operations during processing of cereal-based products on the levels of deoxynivalenol, total aflatoxin, ochratoxin A, and zearalenone: A systematic review and meta-analysis. *Food Chemistry* 268: 611–624. <https://doi.org/10.1016/j.foodchem.2018.06.072>
- Nabizadeh, S., Shariatifar, N., Shokoohi, E., Shoeibi, S., Gavahian, M., Fakhri, Y., Azari, A. and Khaneghah, A.M., 2018. Prevalence and probabilistic health risk assessment of aflatoxins B₁, B₂, G₁, and G₂ in Iranian edible oils. *Environmental Science and Pollution Research* 25: 35562–35570. <https://doi.org/10.1007/s11356-018-3510-0>
- Ostadrhimi, A., Ashrafnejad, F., Kazemi, A., Sargheini, N., Mahdavi, R., Farshchian, M. and Mahluji, S., 2014. Aflatoxin in raw and salt-roasted nuts (pistachios, peanuts and walnuts) sold in markets of Tabriz, Iran. *Jundishapur Journal of Microbiology* 7: 1–4. <https://doi.org/10.5812/jjm.8674>
- Pak-kish, Z. and Rahemi, M., 2005. The effect of gibberellic acid on reduced cracking of pistachio hull, aflatoxin, and quality of dried pistachio fruit. *Journal of Science and Technology of Agriculture and Natural Resources* 9: 69–78. [In Persian]
- Panahirad, S., Zaare-Nahandi, F., Mohammadi, N., Alizadeh-Salteh, S. and Safaie, N., 2014. Effects of salicylic acid on *Aspergillus flavus* infection and aflatoxin B₁ accumulation in pistachio (*Pistacia vera* L.) fruit. *Journal of the Science of Food and Agriculture* 94: 1758–1763. <https://doi.org/10.1002/jsfa.6488>
- Pascale, M.N., 2009. Detection methods for mycotoxins in cereal grains and cereal products. *Zbornik Matice srpske za prirodne nauke* 117: 15–25. <https://doi.org/10.2298/ZMSPN0917015P>
- Pereira, V., Fernandes, J. and Cunha, S., 2014. Mycotoxins in cereals and related foodstuffs: A review on occurrence and recent methods of analysis. *Trends in Food Science & Technology* 36: 96–136. <https://doi.org/10.1016/j.tifs.2014.01.005>

- Piğłowski, M., 2019. Comparative analysis of notifications regarding mycotoxins in the Rapid Alert System for Food and Feed (RASFF). *Quality Assurance and Safety of Crops & Foods* 11: 725–735. <https://doi.org/10.3920/QAS2018.1398>
- Rahaie, S., Emam-Djomeh, Z., Razavi, S. and Mazaheri, M., 2010a. Immobilized *Saccharomyces cerevisiae* as a potential aflatoxin decontaminating agent in pistachio nuts. *Brazilian Journal of Microbiology* 41: 82–90. <https://doi.org/10.1590/S1517-83822010000100014>
- Rahaie, S., Razavi, S.H. and Emam Jomeh, Z., 2010b. The ability of *Saccharomyces Cerevisiae* strain in Aflatoxin reduction in Pistachio nuts. *Journal of Food Science and Technology* 7: 81–88. [In Persian]
- Rahmani, J., Alipour, S., Miri, A., Fakhri, Y., Riahi, S.-M., Keramati, H., Moradi, M., Amanidaz, N., Pouya, R.H., Bahmani, Z. and Mousavi Khaneghah, A., 2018. The prevalence of aflatoxin M1 in milk of Middle East region: A systematic review, meta-analysis and probabilistic health risk assessment. *Food and Chemical Toxicology* 118: 653–666. <https://doi.org/10.1016/j.fct.2018.06.016>
- RASFF, 2016. The Rapid Alert System for Food and Feed — Annual report. Available at: https://ec.europa.eu/food/safety/rasff_en (Accessed November 2017).
- Rastegar, H., Shoeibi, S., Yazdanpanah, H., Amirahmadi, M., Khaneghah, A.M., Campagnollo, F.B. and Sant'Ana, A.S., 2017. Removal of aflatoxin B1 by roasting with lemon juice and/or citric acid in contaminated pistachio nuts. *Food Control* 71: 279–284. <https://doi.org/10.1016/j.foodcont.2016.06.045>
- Ros, E., 2010. Health benefits of nut consumption. *Nutrients* 2: 652–682. <https://doi.org/10.3390/nu2070652>
- Rossi, C.N., Takabayashi, C.R., Ono, M.A., Saito, G.H., Itano, E.N., Kawamura, O., Hirooka, E.Y. and Ono, E.Y.S., 2012. Immunoassay based on monoclonal antibody for aflatoxin detection in poultry feed. *Food Chemistry* 132: 2211–2216. <https://doi.org/10.1016/j.foodchem.2011.12.067>
- Sakamoto, S., Putalun, W., Vimolmangkang, S., Phoolcharoen, W., Shoyama, Y., Tanaka, H. and Morimoto, S., 2018. Enzyme-linked immunosorbent assay for the quantitative/qualitative analysis of plant secondary metabolites. *Journal of Natural Medicines* 72: 32–42. <https://doi.org/10.1007/s11418-017-1144-z>
- Serra, M.S., Pulles, M.B., Mayanquer, F.T., Vallejo, M.C., Rosero, M.I., Ortega, J.M. and Naranjo, L.N., 2018. Evaluation of the use of gamma radiation for reduction of aflatoxin B1 in corn (zea mays) used in the production of feed for broiler chickens. *Journal of Agricultural Chemistry and Environment* 7: 21. <https://doi.org/10.4236/jacen.2018.71003>
- Shakeri, P. and Fazaeli, H., 2016. Microbial contamination and concentration of aflatoxin in pistachio by-product. *Journal of Food Microbiology* 3: 61–72. [In Persian]
- Siahmoshteh, F., Siciliano, I., Banani, H., Hamidi-Esfahani, Z., Razzaghi-Abyaneh, M., Gullino, M.L. and Spadaro, D., 2017. Efficacy of *Bacillus subtilis* and *Bacillus amyloliquefaciens* in the control of *Aspergillus parasiticus* growth and aflatoxins production on pistachio. *International Journal of Food Microbiology* 254: 47–53. <https://doi.org/10.1016/j.ijfoodmicro.2017.05.011>
- Suzuki, T., Munakata, Y., Morita, K., Shinoda, T. and Ueda, H., 2007. Sensitive detection of estrogenic mycotoxin zearalenone by open sandwich immunoassay. *Analytical Sciences* 23: 65–70. <https://doi.org/10.2116/analsci.23.65>
- Taghizadeh, S.F., Rezaee, R., Davarynejad, G., Asili, J., Nemati, S.H., Goumenou, M., Tsakiris, I., Tsatsakis, A.M., Shirani, K. and Karimi, G., 2018. Risk assessment of exposure to aflatoxin B1 and ochratoxin A through consumption of different Pistachio (*Pistacia vera* L.) cultivars collected from four geographical regions of Iran. *Environmental Toxicology and Pharmacology* 61: 61–66. <https://doi.org/10.1016/j.etap.2018.05.010>
- Tavakolipour, H., Javanmard Dakheli, M. and Zirjani, L., 2010. The inhibitory effect of edible coating of pistachio kernel basis of whey protein concentrate and Shirazi thyme extract on the production of aflatoxin toxin. *Journal of Innovation in Food Science and Technology* 2: 53–63. [In Persian]
- Tavakolipour, H., Javanmard, M. and Zirjany, L., 2012. Antiaflatoxic activity of pistachio kernel coated by whey protein based edible film incorporated with *zataria multiflora* essential oil. *Journal of Food Science and Technology* 36: 11–19. [In Persian]
- Tavakolipour, H. and Mokhtarian, M., 2014. Estimation of equilibrium moisture content of pistachio powder through the ANN and GA approaches. *International Journal of Food Engineering* 10: 747–755. <https://doi.org/10.1515/ijfe-2013-0022>
- Tavakolipour, H. and Mokhtarian, M., 2016. New approach to the calculation of pistachio powder hysteresis. *International Agrophysics* 30: 245–251. <https://doi.org/10.1515/intag-2015-0093>
- Tavakolipour, H., Mokhtarian, M. and Kalbasi-Ashtari, A., 2017. Lipid oxidation kinetics of pistachio powder during different storage conditions. *Journal of Food Process Engineering* 40: e12423. <https://doi.org/10.1111/jfpe.12423>
- Turner, N.W., Bramhmbhatt, H., Szabo-Vezse, M., Poma, A., Coker, R. and Piletsky, S.A., 2015. Analytical methods for determination of mycotoxins: An update (2009–2014). *Analytica Chimica Acta* 901: 12–33. <https://doi.org/10.1016/j.aca.2015.10.013>
- Yazdanpanah, H., Mohammadi, T., Abouhossain, G. and Cheraghali, A.M., 2005. Effect of roasting on degradation of aflatoxins in contaminated pistachio nuts. *Food and Chemical Toxicology* 43: 1135–1139. <https://doi.org/10.1016/j.fct.2005.03.004>
- Zhang, L., Dou, X.-W., Zhang, C., Logrieco, A.F. and Yang, M.-H., 2018. A review of current methods for analysis of mycotoxins in herbal medicines. *Toxins* 10: 65. <https://doi.org/10.3390/toxins10020065>